

RESEARCH ARTICLE

Open Access

Molecular cloning and tissue expression of the fatty acid-binding protein (*Es-FABP*) gene in female Chinese mitten crab (*Eriocheir sinensis*)

Ya-Nan Gong[†], Wei-Wei Li[†], Jiang-Ling Sun, Fei Ren, Lin He, Hui Jiang, Qun Wang^{*}

Abstract

Background: Fatty acid-binding proteins (FABPs), small cytosolic proteins that function in the uptake and utilization of fatty acids, have been extensively studied in higher vertebrates while invertebrates have received little attention despite similar nutritional requirements during periods of reproductive activity.

Results: Therefore, a cDNA encoding *Eriocheir sinensis* FABP (*Es-FABP*) was cloned based upon EST analysis of a hepatopancreas cDNA library. The full length cDNA was 750 bp and encoded a 131 aa polypeptide that was highly homologous to related genes reported in shrimp. The 9108 bp *Es-FABP* gene contained four exons that were interrupted by three introns, a genomic organization common among FABP multigene family members in vertebrates. Gene expression analysis, as determined by RT-PCR, revealed the presence of *Es-FABP* transcripts in hepatopancreas, hemocytes, ovary, gills, muscle, thoracic ganglia, heart, and intestine, but not stomach or eyestalk. Real-time quantitative RT-PCR analysis revealed that *Es-FABP* expression in ovary, hemocytes, and hepatopancreas was dependent on the status of ovarian development, with peak expression observed in January.

Conclusions: Evidence provided in the present report supports a role of *Es-FABP* in lipid transport during the period of rapid ovarian growth in *E. sinensis*, and indirectly confirms the participation of the hepatopancreas, ovary, and hemocytes in lipid nutrient absorption and utilization processes.

Background

Fatty acid-binding proteins (FABPs) are small (14-15 kDa) cytosolic proteins that bind non-covalently to hydrophobic ligands, primarily fatty acids [1]. Physiological roles of FABP include, but are not limited to, the uptake and utilization of fatty acids, intracellular targeting of fatty acids to specific organelles and metabolic pathways, and the protection of cellular structures from the detergent effects of fatty acids [2-4]. Similarity among FABP deduced amino acid sequences in vertebrates and invertebrates are generally low despite the highly conserved gene structure of four exons and three introns of variable size [5,6], with the exception of desert locust FABP3 [7] and zebrafish FABP1a [8]. Tertiary structure is common among all FABP family members, and consists of ten antiparallel β -sheet strands that surround the ligand binding domain [9].

To date, 12 FABP isoforms have been identified in vertebrates [10]. While genes were originally named according to the initial tissue from which they were first isolated, *e.g.* Liver-type FABP (L-FABP), Intestinal-type FABP (I-FABP), Heart-type FABP (H-FABP), Adipocyte-type FABP (A-FABP), Epidermal-type (E-FABP), isoform expression among multiple tissues and differences in tissue distribution among FABP orthologs have resulted in the implementation of numeric nomenclature, such that FABP1 corresponds to L-FABP, FABP2 to I-FABP, FABP3 to H-FABP, FABP4 to A-FABP, and FABP5 to E-FABP [9,11].

FABPs bind a single ligand molecule [12], with the exception of FABP1 and FABP10, both L-FABPs capable of simultaneously binding two ligand molecules [13]. FABP1 was the first FABP isolated from liver [14], and functions in fatty acid uptake [15] and metabolic pathway allocation in vertebrates [16], lipoprotein production [17], and nuclear delivery of peroxisome proliferator activated receptor (PPAR) ligands that

* Correspondence: qun_300@hotmail.com

[†] Contributed equally

School of Life Science, East China Normal University, Shanghai 200062, China

results in the modulation of targeted gene expression [18]. FABP10 was first isolated from chicken liver [13], and is more similar in sequence to ileal FABPs than mammalian FABP1. To date, the FABP10 gene and protein have only been identified in nonmammalian vertebrates [8].

While FABPs among vertebrates have been studied in detail for more than three decades, with more than 400 identified, only ≈ 40 FABPs have yet been identified in invertebrates [1,9] and no information concerning FABP is available for *Brachyura*. The Chinese mitten crab (*Eriocheir sinensis*), a commercially important species in South-East Asia, is widely farmed in China and has quickly become an important aquaculture species [19] that has been cultured in ponds, reservoirs and lakes since the 1990's [20]. Hepatopancreas is generally regarded as a major lipid storage organ analogous to the fat body in insects and adipose tissue and liver in vertebrates [21]. The hepatopancreas in crab is a midgut diverticulum involved in the synthesis and secretion of digestive enzymes, final food digestion, nutrient absorption, and lipid and carbohydrate metabolism [22]; further, it is also a sensitive indicator of lipid metabolism and nutritional status [23]. Previous studies have shown that energy stored in crab hepatopancreas is in preparation for the significant expenditure required during the early stages of reproduction [24]. In order to elucidate potential functions of the hepatopancreas during reproduction in *E. sinensis*, a nonnormalized hepatopancreatic cDNA library was constructed [25]. EST analysis and subsequent cloning revealed an *Es-FABP* unigene, the first FABP gene identified in *Brachyura*. In the present study, we isolated FABP, for the first time from hepatopancreas of the Chinese mitten crab, investigated seasonal expression with respect to reproductive stage, and explored the relationship between FABP expression level and ovary development in the Chinese mitten crab in order to provide further insights into reproduction, nutrition and development of the Chinese mitten crab for farming industry.

Methods

Tissues preparation

Healthy adult female crabs (*E. sinensis*) were purchased from Tongchuan aquatic product market, Shanghai, China. Crabs were placed in an ice-bath for 1-2 min until lightly anesthetized. Eleven tissues were collected, including hepatopancreas, gills, stomach, intestine, cranial ganglia, thoracic ganglia, hemocyte, heart, muscle, and ovary, frozen immediately in liquid nitrogen, and stored at -80°C until nucleic acid extraction.

Based on the ovarian classification of Xue et al [26], ovarian Stage III-1 (40-50 mm diameter), III-2 (60-90 mm diameter), and IV (92-100 mm diameter) specimens

were collected from August to September, September to November, and December to March, respectively, for ovarian *Es-FABP* expression analysis. For the present study, ovarian stages were further sub-divided based upon the month of dissection: Stage III-1 (Aug), Stage III-1 (Sep); Stage III-2 (Sep), Stage III-2 (Oct), Stage III-2 (Nov); Stage IV (Jan). Collection of hepatopancreas and hemocyte specimens occurred concomitantly with ovarian staged specimens. Tissues were immediately frozen in liquid nitrogen and stored at -80°C until RNA extraction.

Nucleic acid extraction

RNA was extracted using a RNA Extraction Kit (Axygen, USA) according to the manufacturer's protocol. RNA concentration and quality were estimated by spectrophotometry at an absorbance of 260 nm, (Eppendorf Biophotometer plus, Germany) and agarose gel electrophoresis (Bio-Rad PowerPacBasic, USA), respectively. Total RNA (200 ng) was reverse transcribed using the PrimeScript™ RT-PCR Kit (TaKaRa, Japan) for semi-quantitative RT-PCR or the SYBR PrimeScript™ RT-PCR Kit (TaKaRa, Japan) for real-time quantitative RT-PCR (qRT-PCR).

Chinese mitten crab genomic DNA was extracted from hepatopancreas using the Axyprep™ Multisource Genomic DNA Miniprep Kit (Axygen, USA) according to manufacturer's protocol. DNA concentration and quality was estimated by spectrophotometry at an absorbance of 260 nm (Eppendorf Biophotometer plus, Germany) and agarose gel electrophoresis (Bio-Rad PowerPacBasic, USA), respectively.

EST analysis and full-length cDNA cloning

A cDNA library was previously constructed from the hepatopancreas of Chinese mitten crab, and 3355 successful sequencing reactions were obtained using a T3 primer [25]. BLASTx analysis of EST sequences revealed 20 ESTs with high identity to *FcFABP* in *Fenneropenaeus chinensis* (ACU82845), which could be assembled into a single contig [a representative 527 bp EST is reported as GenBank: FG358115]. Sequence of the assembled *Es-FABP* contig was confirmed from the 3' direction using a T7 primer (Table 1).

To verify cDNA sequence, a gene-specific primer pair, F-S2 (sense) and F-R2 (anti-sense) (Table 1), was designed by Primer Premier 5.0 based on the sequence of the assembled contig mentioned above. The PCR reaction was performed in a ABI 2720 Thermal Cycler in a total volume of 25 μl and contained 2.5 μl of $10 \times$ PCR buffer (Mg^{2+} plus), 2.0 μl of dNTPs mix (2.5 mM each), 0.5 μl of each primer (20 μM), 18.375 μl of RNase-free water, 0.125 μl (5 U/ μl) of Taq polymerase (TaKaRa, Japan), and 1 μl cDNA (500 ng/ μl) as

Table 1 Oligonucleotide primers used to amplify the *E. sinensis* FABP gene

Primer name	Sequence (5'-3')	Application
T7	TAATACGACTCACTATAGG	cDNA cloning
SP6	ATTTAGGTGACACTATAGAA	cDNA cloning
5'SP1	CCCACAGCACTGAGCCCAATC	Genome walking
5'SP2	TATTTCCCGGTGATGGACATGA	Genome walking
5'SP3	GGTAGAAACAGCGAGGGCAACT	Genome walking
3'SP1	CACCTCCGCCAAGATTGTTATT	Genome walking
3'SP2	CGCACTTGAAGAACCATAGCAG	Genome walking
3'SP3	GCTTTATGAGATTCGTTTGC GTGAT	Genome walking
F-R2	CAGAAGATGTTACAAGACTAAAG	RT-PCR
F-S2	GAGITGCCCTCGCTGTTTGCTAT	RT-PCR
F-R3	CGTGTCTTGATGACGATGT	RealTime-PCR
F-S3	TGGCTCAGTGCTGTGGGGT	RealTime-PCR
Actin-R	CTCCTGCTTGCTGATCCACATC	RT & RealTime-PCR
Actin-S	GCATCCACGAGACCACTTACA	RT & RealTime-PCR

template. The PCR conditions were as follows: 94°C for 5 min; 30 cycles of 94°C for 30 s, 58°C for 30 s, 72°C for 30 s; and 72°C for 10 min. Appropriately sized PCR products were gel-purified and ligated into a pGEM-T easy vector (Promega, USA) with T4 DNA ligase. Positive clones containing inserts of predicted size were sequenced using T7 and SP6 primers (Table 1).

Cloning of the *Es-FABP* gene and associated 5', 3'-flanking regions

The *Es-FABP* genomic DNA sequence was obtained with a gene-specific primer pair (F-S1, F-R1; Table 1), which corresponded to the cloned cDNA sequence. The final PCR reaction was performed in a total volume of 25 µl as described above. The 5' and 3'-flanking regions were cloned using a Genome Walking Kit (TaKaRa, Japan) according to the manufacturer's protocol with the following primers: 5'SP1, 5'SP2, 5'SP3, 3'SP1, 3'SP2, and 3'SP3. Specific products were purified and ligated into a pGEM-T easy vector (Promega, USA) with T4 DNA ligase. Positive clones were sequenced using SP6 primers (Table 1) and a 3730XL DNA analyzer (Applied Biosystems, Foster City, CA). The positions of exons and introns were determined by spidey. <http://www.ncbi.nlm.nih.gov/IEB/Research/Ostell/Spidey/>

Phylogenetic analysis

Es-FABP nucleotide and deduced amino acid sequences were compared to those reported for other organisms using the BLAST algorithm at the National Center for Biotechnology Information <http://www.ncbi.nlm.nih.gov/> to assess sequence identity. FABP amino acid sequences from *E. sinensis* and representative taxa were retrieved from NCBI GenBank and analyzed using the ClustalW Multiple Alignment program <http://www.ebi.ac.uk/>

clustalw/. The open reading frame (ORF) of the cloned *Es-FABP* cDNA was determined by the ORF Finder <http://www.ncbi.nlm.nih.gov/gorf/>. A neighbor-joining (NJ) phylogenetic tree was constructed using the MEGA software version 4.0 package (<http://www.megasoftware.net/>). The confidence values for phylogeny analysis were determined by 1000 bootstrap replicates and expressed as a percentage. Three-dimensional domain structure of *Es-FABP* was predicted by Protein existence Server. http://www.uniprot.org/manual/protein_existence.

Semi-quantitative RT-PCR analysis

Total RNA was extracted and reverse transcribed as described above. A predicted FABP PCR amplicon of 500 bp was generated using the gene-specific primer pair F-R2 and F-S2 (Table 1). The constitutively expressed beta-actin (Actin-R and Actin-S) produced a 276 bp product and served as an internal control. FABP and beta-actin genes were run in two single PCR reaction, with same PCR conditions and cycle number. The PCR reaction was performed in a final volume of 25 µl and contained 2.5 µl 10 × PCR buffer (Mg²⁺ Plus), 2 µl 10 mM dNTP mixture, 0.25 µl 20 mM each primer, 18.875 µl PCR-Grade water, 0.125 µl 5 U Ex Taq™ Hot Start Version (TaKaRa, Japan), and 1 µl cDNA as template. PCR conditions were as follows: 94°C for 5 min; 30 cycles of 94 °C for 30 s, 55°C for 30 s, 72°C for 1 min; and 72°C for 5 min. RT-PCR products were separated on 1.5% agarose gel with ethidium bromide and detected under a Gel Doc 2000 ultraviolet light (Tannon, China).

SYBR Green real-time qRT-PCR analysis

Real-time qRT-PCR was performed in a C1000™ Thermal Cycler (BioRad CFX 96™ Real-Time System) according to the manufacturer's instructions. PCR conditions were as follows: 95°C for 30 s; 40 cycles of 95°C for 5 s, 60°C for 30 s, with an 0.1°C/s incremental increase from 60°C to 95°C. The final volume of each qRT-PCR reaction was 25.0 µl, and contained 12.5 µl 2 × SYBR Premix Ex Taq (TaKaRa, Japan), 0.5 µl diluted cDNA template, 11.0 µl PCR-Grade water, and 0.5 µl of each 20 uM primer (F-R3, F-S3 Table 1). Beta-actin fragments were amplified using the primer pair Actin-R and Actin-S, and served as a positive control. Samples were run in triplicate, and *Es-FABP* expression levels were calculated by the 2^{-ΔΔCt} comparative CT method [27]. Data are represented as triplicate mean ± SD (standard deviation) and presented as the n-fold difference relative to beta-actin.

Statistical analysis

Statistical significance was determined by a one-way ANOVA and a posthoc Tukey test using SPSS 11.5 software. Significance was set at *P* < 0.05.

Results

Cloning and identification of the *Es-FABP* cDNA

A full length FABP cDNA (*Es-FABP*) was isolated from the hepatopancreas of a female Chinese mitten crab [GenBank: GU568242]. The full length cDNA (750 bp) contained a 393 bp open reading frame (ORF), which encoded a putative 131 amino acid FABP protein, a 65 bp 5'-untranslated region (UTR), and a 282 bp 3'-UTR (Fig. 1). A single polyadenylation signal (AATAA) was observed 719 bp upstream of the 12 bp poly (A) tail. Analysis of the *Es-FABP* cDNA

sequence using BLASTX revealed significant sequence similarity to other FABPs sequences included in the National Center for Biotechnology Information database (NCBI, <http://www.ncbi.nlm.nih.gov/BLAST/>).

Genomic sequence and structure of *Es-FABP*

The *Es-FABP* gene was 9108 bp in length and was comprised of four exons interrupted by three introns (Fig. 2) [GenBank: GU568243]. Typical intron-exon junction structures with donor and acceptor (GT and AG)


```
1          attccgaattcggagttgcc
21 ctcgctgtttctaccaacaccaccaccaacaacaacaccatc
66 atgtccatcaccgggaatacgttctgagcggcaacgagaaatat
   M S I T G K Y V L S G N E K Y 15
111 gctgattggctcagtgetgtgggtatccccgctgacctcgccgcc
   A D W L S A V G I P A D L A A 30
156 aagctggaggccgccaagccctccctggatgtgaccagagtggc
   K L E A A K P S L D V T Q S G 45
201 aacgacatcgtcatcaagaccacggccggcgacaagaacttcacc
   N D I V I K T T A G D K N F T 60
246 aacaccatcacgctcggcaaggagtcacaggccacgctccccggc
   N T I T L G K E S Q A T L P G 75
291 ggcacatcgagtacacgggtgaagttgacgctctctggcaccaccctc
   G I E Y T V K L T L S G T T L 90
336 accggcacctgggactttggcggcaagactggcaccgctaccgtc
   T G T W D F G G K T G T A T V 105
381 gagttcaactgccgacggggtcacacagaccatgatctacaacaac
   E F T A D G V T Q T M I Y N N 120
426 atcacctccaagaggatctactcccgccagtagtctgctacaccc
   I T S K R I Y S R Q *          131
471 caagtctatcaatatggtctctagattaccttttcctcacttcac
516 ttgctaaatttggttatctttgctcttcagcctgacttagtaact
561 ttggtatttaattgttacaccacctgcaactcaagtcacttttat
606 ctttacattcttttagtcttgaacatcttctgaagtgagtaattt
651 agcaaaaagacatccttgaacactatctgtactgctaacaataag
696 gatgtgaaattttatttctaataatcattaaagctttttttataa
   aaaaaaaaaa
```

Figure 1 *Es-FABP* nucleotide (above) and deduced amino acid (below) sequences. Nucleotides numbering was initiated at the first base pair at the 5' end. Amino acid numbering began with the first in-frame methionine. The stop codon is marked by an asterisk. The polyadenylation signal (AATAA) is enclosed by a black ellipse. The *Es-FABP* sequence was submitted to GenBank [GenBank: GU568242].



dinucleotide sequences were conserved in *Es-FABP*. *Es-FABP* genomic and cDNA sequences exhibited 99% similarity. The 1201 bp 5' flanking upstream region contained a putative TATA box, but no binding motifs were identified. Comparison of the exon/intron

organization of the *FABP* genes from *Danio rerio* (*fabp10*), *Gallus gallus* (*fabp10*), *Ciona intestinalis* (*fabp10*), *Homo sapiens* (*fabp1*), *Mus musculus* (*fabp1*) (Fig. 3). The positions of exons and introns were determined by an mRNA-to-genomic alignment using Spidey.



Homology Analysis of Es-FABP

The identity of Es-FABP and other representative vertebrate and invertebrate deduced amino acid FABP sequences was explored via multiple sequence alignment using ClustalX (Fig. 4) [28]. Overall percent identity was high among Es-FABP and other reported invertebrate sequences, with 56% identity to *Fenneropenaeus chinensis* and 62% identity to *Litopenaeus vannamei*. Overall percent identity of Es-FABP and representative higher and lower vertebrates was substantially lower, with 21% identity to *Ctenopharyngodon idella*; 22% identity to *Sparus aurata*; 27% identity to *Xenopus tropicalis*; 28% identity to *Rattus norvegicus*; 29% identity to *Homo sapiens*, *Mus musculus*, *Danio rerio*, *Taeniopygia guttata*, *Rana catesbeiana*, *Pongo abelii*; 30% identity to *Bos Taurus*; 32% identity to *Gallus gallus*.


A NJ phylogenetic tree was constructed based on reported FABPs amino acid sequences using MEGA4.0 software (Fig. 5). The reliability of the branching was tested using the bootstrap resampling (with 1000 pseudo

replicates) technique. Two distinct sister groups were observed, with tree topology in agreement with traditional taxonomic relationships. The first group contained vertebrate FABP1, FABP6 and FABP10 sequences, as well as Es-FABP and shrimp FABPs. The second group contained vertebrate FABP2, FABP3, FABP4, FABP5, FABP7, FABP8, FABP9, FABP11, and FABP12 sequences.

A homology model of Es-FABP was predicted in SWISS-MODEL database <http://swissmodel.expasy.org/> revealed conservation of tertiary structure, with the 10 anti-parallel β -strands forming a barrel and a clamshell-like structure.

Tissue distribution and reproductive dependence of Es-FABP expression

Semi-quantitative RT-PCR and real-time qRT-PCR were employed to investigate the distribution of *Es-FABP* mRNA in different tissues as well as to assess expression during the seasonal female reproductive cycle. As



determined by RT-PCR, *Es-FABP* was widely distributed with high and detectable expression levels observed in hepatopancreas, ovary, hemocytes, and gills, muscle, cranial ganglia, thoracic ganglia, heart, intestine, respectively; while expression was nondetectable in stomach and eyestalk (Fig. 6).

As determined by real-time qRT-PCR, beta-actin normalized *Es-FABP* expression in ovary increased rapidly and significantly from Stage III-1 (Aug-Sep) to reach peak expression during Stage III-2 (Nov) (Fig. 7). *Es-FABP* ovarian expression then decreased significantly through Stage IV (Jan). Hepatopancreatic expression decreased gradually from Stage III-1 (Aug) to Stage III-2 (Nov) and then increased significantly in Stage IV (Jan) (Fig. 8). *Es-FABP* expression in hemocytes did not differ significantly from Stage III-1 (Aug) through Stage III-2 (Nov), with a sharp peak observed during Stage IV (Jan) (Fig. 9).

Discussion

FABP gene family members are evolutionarily conserved, with extensive and interdependent functions in

the regulation of fatty acid oxidation/metabolism [9]. Although much research has focused on the classification, structure and function of vertebrate FABPs, these proteins remain unidentified in many invertebrate species. The current study reports the cloning and gene structure of a FABP family member in *E. sinensis*, *Es-FABP*, and presents evidence of functionality in fatty acid metabolism.

The full-length *Es-FABP* cDNA reported in the present paper encoded a putative FABP protein of 131 amino acids, which falls within the size range of other reported FABPs (127-136 amino acids) [29]. The ClustalX alignment of *Es-FABP* and fourteen other reported vertebrate and invertebrate FABP sequences revealed high identity among invertebrate sequences (56-62%) although percent identity remained lower when comparing within invertebrate and mammalian taxa (79-90%) [9]. Three-dimensional homology modeling revealed that several key tertiary structures of *Es-FABP* were similar to those of vertebrate FABP, such as the 10 anti-parallel β -strands, their resultant barrel with a clam-shell-like structure, and the barrel cap comprised of a







Figure 5 Neighbor-Joining phylogenetic tree of representative vertebrate and invertebrate FABP amino acid sequences. Bootstrap values supporting branch points are expressed as the percentage of 1000 replicates. The following organisms with FABP GenBank reported sequences were included in the analysis: *A. mexicanum* (Am-FABP10) [GenBank: P81400.2]; *B. arenarum* (Ba-FABO10) [P83409]; *D. rerio* (Dr-FABP1b) [AAZ08576], Dr-FABP1a [NP_001038177], Dr-FABP3 [NP_694493], Dr-FABP6 [ACD37360], Dr-FABP10 [AAI64928], Dr-FABP11 [NP_001004682]; *F. chinensis* (Fc-FABP) [ACU82845]; *G. gallus* (Gg-FABP1) [NP_989523], Gg-FABP4 [NP_989621]; *H. sapiens* (Hs-FABP1) [NP_001434], Hs-FABP4 [NM_001442], Hs-FABP5 [NP_001435], Hs-FABP6 [AAH22489], Hs-FABP7 [CAG33338], Hs-FABP8 [NP_002668], Hs-FABP9 [NP_001073995], Hs-FABP12 [NP_001098751]; *L. vannamei* (Lv-FABP10) [ABD65306]; *M. musculus* (Mm-FABP1) [NP_059095], Mm-FABP2 [AAS00550], Mm-FABP5 [NP_034764], Mm-FABP7 [CAJ18607], Mm-FABP9 [NP_035728], Mm-FABP2 [NP_083586]; *O. mykiss* (Om-FABP3) [NP_001118185]; *P. abelii* (Pa-FABP1) [NP_001125017]; *R. norvegicus* (Rn-FABP3) [NP_077076], Rn-FABP6 [NP_058794], Rn-FABP9 [NP_074045], Rn-FABP2 [NP_001128086]; *R. scapo* (Rs-FABP10) [P80856]; *S. scrofa* (Ss-FABP2) [NM_001031780], Ss-FABP4 [NM_001002817], Ss-FABP5 [NP_001034835], Ss-FABP7 [AAY17257], Ss-FABP11 [CAM58515]; *T. guttata* (Tg-FABP1) [XP_002188068]; *X. laevis* (Xi-FABP2) [NM_001085877].



pair of α -helices, which enclose the FABP lipid-binding site cavity [30].


The cloned *Es-FABP* gene was 9108 bp in length, and was comprised of four exons interrupted by three introns, a conserved genomic organization among both species [31,32] and FABP superfamily members. Comparison of the exon/intron organization of the *FABP* genes revealed that, the genes are organized in four exons and three introns located in conserved positions, although varying markedly in size. And the second intron of *Es-FABP* was substantially longer than reported for other species. In general, while large introns are not unusual in organisms with very large genomes, they are less common among heavily transcribed genes, like the locust *FABP* [7]. Long introns might be favored because they enhance recombination, thus introducing mutations in adjacent exons [33] or



intra- and inter- gene recombination [34], all of which would be subjected to selection pressures.

Phylogenetic analysis revealed two distinct sister groups within our constructed NJ tree as previously reported [8,35]: (1) group 1 contained FABP1, 6, 10, and (2) group 2 contained FABP2, 3, 4, 5, 7, 8, 9, 11, 12. Schaap *et al.* estimated that FABP1, FABP6 and FABP10 diverged from a common ancestral gene ~679 million years ago (mya) [6]. FABP6 is an ileal-type FABP [36], while FABP1 and FABP10 are paralogs, and are thus evolutionarily related liver-type FABPs, which possibly arose by a gene duplication event [8]. *Es-FABP* segregated within group 1, clustering in close proximity to Fc-FABP10 and LvFABP10. ClustalX alignment results suggest that *Es-FABP* shares a high percent identity with FABP10. Based on collective phylogenetic evidence, we hypothesize that the *Es-FABP* we cloned from hepatopancreas is a FABP10.

We hypothesized that *Es-FABP* tissue expression profiles may provide useful cues when speculating gene function. In the present study, *Es-FABP* transcripts were primarily detected in hepatopancreas, hemocytes, and ovary. The hepatopancreas is generally regarded as a major storage organ of lipids, which are transported to reproductive organs and tissues during a period of reproductive activity [24]. Interestingly, the crustacean ovary contains a high lipid content compared to other organs [37,38], which is a key nutrient required for ovarian development and appropriate egg hatching rate and larval survival [38-41]. Further, the crustacean ovary is rich in long-chain poly unsaturated fatty acids (PUFA) [42]. As intracellular transporters of PUFA, FABP may serve as mediators of their physiological function, availability and access to intracellular targeted systems [10].



The high *Es-FABP* expression levels observed in these organs suggest that lipid nutrients, especially fatty acids, are transported from the hepatopancreas to the ovary in great supply via *Es-FABP* expressing hemocytes during the stage of rapid ovarian development. *Es-FABP* transported lipids are then stored in the ovaries in preparation for the significant and impendent energy expenditure required during reproduction [21].

Real-time qRT-PCR analysis revealed *Es-FABP* expression levels were dependent on the stage of ovarian development in all three organs. Fatty acids (FA) in general, including the ubiquitous palmitic and oleic acids, can regulate gene expression at the transcriptional level [43,44]. Furthermore, long-chain FA and long-chain dicarboxylic acids can act as inducers of *FABP1* expression [45,46], as tissue FABP content is related to the rate of FA uptake and/or utilization [47-49]. Similarly, manipulations altering the rate of tissue FA metabolism appear to be associated with concomitant changes in FABP content [50]. Therefore, a direct or indirect feedback loop between FA and FABP may exist. In the present study, ovarian *Es-FABP* expression increased with the developmental progression of the ovary, mirroring lipid nutritional requirements. The ovary requires substantial accumulation of lipids, especially fatty acids, during developmental stages [51-53]. Interestingly, the

rapid increase in ovarian lipid deposition was associated with a gradual decline in *Es-FABP* expression in hepatopancreas, suggesting hepatopancreas-mediated lipid transport may be occurring during ovarian development. Determining the existence or absence of a hepatopancreas-ovary feedback loop requires further study.

Conclusions

In conclusion, evidence provided in the present report supports a role of *Es-FABP* in lipid transport during the period of rapid ovarian growth in *E. sinensis*, and indirectly confirms the participation of the hepatopancreas, ovary, and hemocytes in lipid nutrient absorption and utilization processes. However, further study concerning *Es-FABP*'s specific mechanistic role in the process of lipid transport is both required and warranted.

Acknowledgements

This research was supported by the National Natural Science Foundation of China (Grant No. 30671607, 30972241), program of Shanghai Education Commission (No. 08SG24).

Authors' contributions

YNG carried out the molecular studies, participated in the gene cloning, sequence alignment and drafted the manuscript. WWL participated in the phylogenetic analysis, semi-quantitative RT-PCR and SYBR Green real-time qRT-PCR analysis. JLS participated in the tissue preparation. FR participated in the nucleic acid extraction. LH participated in the statistical analysis. HJ

provided EST sequences. QW conceived of the study, and participated in its design and coordination. All authors read and approved the final manuscript.

Received: 10 April 2010 Accepted: 16 September 2010
Published: 16 September 2010

References

- Esteves A, Ehrlich R: Invertebrate intracellular fatty acid binding proteins. *Comp Biochem Physiol C* 2006, **2**:262-74.
- Corsico B, Liou HL, Storch J: The alpha-helical domain of liver fatty acid binding protein is responsible for the diffusion-mediated transfer of fatty acids to phospholipid membranes. *Biochemistry* 2004, **43**:3600-3607.
- Ho SY, Delgado L, Storch J: Monoacylglycerol metabolism in human intestinal Caco-2 cells: evidence for metabolic compartmentation and hydrolysis. *Biol Chem* 2002, **277**:1816-1823.
- Storch J, Veerkamp JH, Hsu KT: Similar mechanisms of fatty acid transfer from human and rodent fatty acid-binding proteins to membranes: liver, intestine, heart muscle and adipose tissue FABPs. *Mol Cell Biochem* 2002, **239**:25-33.
- Bernlohr DA, Simpson MA, Hertzell AV, Banaszak LJ: Intracellular lipid-binding proteins and their genes. *Annu Rev Nutr* 1997, **17**:277-303.
- Schaap FG, Vusse GJ, Glatz JF: Evolution of the family of intracellular lipid binding proteins in vertebrates. *Mol Cell Biochem* 2002, **239**:69-77.
- Wu QW, Andolfatto P: Cloning and sequence of the gene encoding the muscle fatty acid binding protein from the desert locust, *Schistocerca gregaria*. *Ins Biochem Mol Biol* 2001, **31**:553-562.
- Sharma MK, Liu RZ, Thisse C, Thisse B: Hierarchical subfunctionalization of FABP1a, FABP1b and FABP10 tissue-specific expression may account for retention of these duplicated genes in the zebrafish (*Danio rerio*) genome. *FEBS J* 2006, **273**:3216-3229.
- Zimmerman AW, Veerkamp JH: New insights into the structure and function of fatty acid-binding proteins. *Cell Mol Life Sci* 2002, **11**:1096-1116.
- Liu RZ, Li X, Godbout R: A novel fatty acid-binding protein (FABP) gene resulting from tandem gene duplication in mammals: transcription in rat retina and testis. *Genomics* 2008, **92**:436-445.
- Hertzell AV, Bernlohr DA: The mammalian fatty acid-binding protein multigene family: molecular and genetic insights into function. *Trends Endocrin Metab* 2000, **11**:175-180.
- Nichesola D, Perduca M, Capaldi S, Carriso ME, Righetti PG, Monaco HL: Crystal structure of chicken liver basic fatty acid-binding protein complexed with cholic acid. *Biochemistry* 2004, **43**:14072-14079.
- Ceciliani F, Monaco HL, Ronchi S, Faotto L, Spadon P: The primary structure of a basic (pI 9.0) fatty acid-binding protein from liver of *Gallus domesticus*. *Comp Biochem Physiol B* 1994, **109**:261-71.
- Schleicher CH, Santome JA: Purification, characterization, and partial amino acid sequencing of an amphibian liver fatty acid binding protein. *Biochem Cell Biol* 1996, **74**:109-115.
- Newberry EP, Xie Y, Kennedy S, Han X, Buhman KK, Luo J, Gross RW, Davidson NO: Decreased hepatic triglyceride accumulation and altered fatty acid uptake in mice with deletion of the liver fatty acid-binding protein gene. *J Biol Chem* 2003, **278**:51664-51672.
- Storch J, Corsico B: The Emerging Functions and Mechanisms of Mammalian Fatty Acid-Binding Proteins. *Annu Rev Nutr* 2008, **28**:73-95.
- Spann NJ, Kang S, Li AC, Chen AZ, Newberry EP, Davidson NO, Hui STY, Davis RA: Coordinate transcriptional repression of liver fatty acid-binding protein and microsomal triglyceride transfer protein blocks hepatic very low density lipoprotein secretion without hepatosteatosis. *J Biol Chem* 2006, **281**:33066-33077.
- Wolfrum C, Borrmann CM, Borchers T, Spener F: Fatty acids and hypolipidemic drugs regulate peroxisome proliferator-activated receptors α - and γ -mediated gene expression via liver fatty acid binding protein: A signaling path to the nucleus. *Biol Sci Biochem* 2001, **98**:2323-2328.
- Yang WL, Zhang GH: Current trends of aquaculture production and sustainable development of Chinese mitten crab, *Eriocheir sinensis*. *Chin J Freshwater Fish* 2005, **35**:62-64.
- Chen DW, Zhang M, Shrestha S: Compositional characteristics and nutritional quality of Chinese mitten crab (*Eriocheir sinensis*). *Food Chem* 2007, **103**:1343-1349.
- Wen XB, Chen LQ, Ai CX, Zhou ZL, Jiang HB: Variation in lipid composition of Chinese mitten-handed crab, *Eriocheir sinensis* during ovarian maturation. *Comp Biochem Physiol B* 2001, **130**:95-104.
- Verri T, Mandal A, Zilli L, Bossa D, Mandal PK, Ingrosso L, Zonno V, Vilella S, Ahearn GA, Storelli C: D-glucose transport in decapod crustacean hepatopancreas. *Comp Biochem Physiol A* 2001, **130**:585-606.
- Li E, Chen L, Zeng C, Yu N, Xiong Z, Chen X, Qin JG: Comparison of digestive and antioxidant enzymes activities, haemolymph oxyhemocyanin contents and hepatopancreas histology of white shrimp, *Litopenaeus vannamei*, at various salinities. *Aquaculture* 2008, **274**:80-86.
- Normant M, Chrobak M, Szaniawska A: Energy value and chemical composition (CHN) of the Chinese mitten crab *Eriocheir sinensis* (Decapoda: Grapsidae) from the Baltic Sea. *Thermochimica Acta* 2002, **394**:233-237.
- Jiang H, Cai YM, Chen LQ, Zhang XW, Hu SN, Wang Q: Functional annotation and analysis of expressed sequence tags from the hepatopancreas of mitten crab (*Eriocheir sinensis*). *Mar Biotechnol* 2009, **11**:317-326.
- Xue LZ, Du NS, Lai W: Histology of female reproductive system in Chinese mitten-handed crab, *Eriocheir sinensis* (Crustacean, Decapoda). *J East China Norm Univ Nat Sci Ed* 1987, **3**:88-97.
- Livak KJ, Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2^{-Delta Delta C} (T) Method. *Methods* 2001, **25**:402-408.
- Thompson JD, Higgins DJ, Gibson TJ: CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucl Acids Res* 1994, **22**:4673-4680.
- Chen ZH, Xu LM, Shan AS: Fatty acid-binding proteins (FABPs) and their genes of FABPs. *J NE Agr Univ* 2006, **5**:689-692.
- Chmurzynska A: The multigene family of fatty acid-binding proteins (FABPs): function, structure, and polymorphism. *J Appl Genet* 2006, **47**:39-48.
- Hauerland NH: Fatty acid binding proteins in locust and mammalian muscle. Comparison of structure, function, and regulation. *Comp Biochem Physiol B* 1994, **109**:199-208.
- Veerkamp JH, Maatman R: Cytoplasmic fatty acid-binding proteins: their structure and genes. *Prog Lipid Res* 1995, **34**:17-52.
- Cameron JM, Kreitman M: The correlation between intron length and recombination in *Drosophila*. *Dynamic equilibrium between mutational and selective forces*. *Genetics* 2000, **156**:1175-1190.
- Duret L: Why Do Genes Have Introns? Recombination Might Add a New Piece to the Puzzle. *Trends Genet* 2001, **17**:172-175.
- Ananda BV, Christine T, Bernard T, Jonathan MW: Differential tissue-specific distribution of transcripts for the duplicated fatty acid-binding protein 10 (FABP10) genes in embryos, larvae and adult zebrafish (*Danio rerio*). *FEBS J* 2009, **276**:6787-6797.
- Schleicher CH, Cordoba OL, Santome JA: Molecular evolution of the multigene family of intracellular lipid-binding proteins. *Biochem Mol Biol Int* 1995, **36**:1117-1125.
- Ando T, Kanazawa A, Teshima S: Variation in the lipids of tissues during the molting cycle of prawn. *Bull Jpn Soc Sci Fish* 1977, **43**:1445-1449.
- Teshima S, Kanazawa A: Variation in lipid compositions during the ovarian maturation of the prawn. *Bull Jpn Soc Sci Fish* 1983, **49**:957-962.
- Xu XL, Ji WJ, Castell JD: Essential fatty acid requirement of the Chinese prawn, *Penaeus chinensis*. *Aquaculture* 1994, **127**:29-40.
- Lautier J, Lagarrigue JG: Lipid metabolism of the crab *Pachyrapus marmoratus* during vitellogenesis. *Biochem Syst Ecol* 1998, **16**:203-212.
- Alava VR, Kanazawa A, Teshima S, Koshio S: Effects of dietary phospholipids and ω -3-high unsaturated fatty acid on ovarian development of kuruna prawn. *Nippon Suisan Gakk* 1993, **59**:345-351.
- San GJP, Chew EY: The role of omega-3 long-chain polyunsaturated fatty acids in health and disease of the retina. *Prog Retin Eye Res* 2005, **24**:87-138.
- Distel RJ, Robinson GS, Spiegelman BM: Fatty acid regulation of gene expression. Transcriptional and post-transcriptional mechanism. *Biol Chem* 1992, **267**:5937-5941.
- Clarke SD, Jump DB: Regulation of gene transcription by polyunsaturated fatty acids. *Prog Lipid Res* 1993, **32**:139-149.

45. Kaikaus RM, Chan WK, Montellano P, Bass NM: **Mechanisms of regulation of liver fatty acid-binding protein.** *Mol Cell Biochem* 1993, **123**:93-100.
46. Kaikaus RM, Sui Z, Lysenko N: **Regulation of pathways of extramitochondrial fatty acid oxidation and liver fatty acid-binding protein by long-chain monocarboxylic fatty acids in hepatocytes.** *Biol Chem* 1993, **268**:26866-26871.
47. Glatz JFC, Vusse GJ: **Cellular fatty acid-binding proteins: current concepts and future directions.** *Molec Cell Biochem* 1990, **98**:237-251.
48. Ockner RK, Lysenko N, Manning JA, Monroe SE, Barnett DA: **Sex steroid modulation of fatty acid utilization and fatty acid binding protein concentration in rat liver.** *Clin Invest* 1980, **65**:1013-1023.
49. Breda E, Keizer HA, Vork MM, Jong YF, Vusse GJ, Glatz JFC: **Modulation of fatty acid-binding protein content of rat heart and skeletal muscle by endurance training and testosterone treatment.** *Eur J Phys* 1992, **421**:274-279.
50. Glatz JFC, Borchers T, Spener F, Vusse GJ: **Fatty Acids in Cell Signalling: Modulation by Lipid Binding Proteins.** *Prostag Leukotr Ess* 1995, **52**:121-127.
51. Gonzalen-Baro MR, Pollero RJ: **Lipid characterization and distribution in mung tissues of the freshwater crustacean *Macrobrachium borellii* during and annual cycle.** *Comp Biochem Physiol B* 1988, **91**:711-715.
52. Millanena OM: **Effect of fatty acid composition broodstock diet on tissue fatty acid patterns and egg fertilization and hatching in pond reared *Penaeus monodon*.** *Asian Fish* 1989, **2**:127-131.
53. Mourente G, Medina A, Gonzalez S: **Changes in lipid class and fatty acids contents in the ovary and midgut gland of the female fiddler crab *Uca tangeri* (Decapoda, Ocypodiadae) during maturation.** *Mar Biol* 1994, **121**:187-197.

doi:10.1186/1471-2199-11-71

Cite this article as: Gong *et al.*: Molecular cloning and tissue expression of the fatty acid-binding protein (*Es-FABP*) gene in female Chinese mitten crab (*Eriocheir sinensis*). *BMC Molecular Biology* 2010 **11**:71.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

